

Trajectories in quality of life: How to connect to people in the different trajectories?

De Hoogstraat

Content

- What is quality of life?
- Umbrella study and trajectories in quality of life?
- Which factors influence quality of life in persons with SCI?
- Psychosocial rehabilitation in persons with SCI
- Discussion: How can patient organisations and rehabilitation centers work together?



Definition of quality of life

- According to Shalock (2000) there are more than 100 different definitions of QoL.
- According to van Heck (2006) there are as many definitions of QoL as there are scientists in the field of QoL.



How do you define quality of life?

• MD's, PT's, OT's in the UK (N=1572)

– Happiness	72.2%
– Social	25.8%
– Physical	25.2%
– Mental	17.8%
 Independence 	13.0%
 Subjective 	12.3%
– Economic	6.6%
 Good health 	5.3%
– Indefinable	2.0%
– Other	1.6%

(McKevitt, Redfern, et al. Clin Rehabil 2003)



WHO defines quality of life as:


Individuals perception of their position in life **in the context of the culture and value systems** in which they live and in relation to their goals, expectations, standards and concerns





World happiness

A Global Projection of Subjective Well-being: The First Published Map of World Happiness



Happy ----- Average ----- Unhappy



Which factors influence QOL?

- Climate
- Safety
- Education
- Civil status
- Personality
- Social network
- Health (health facilities)
- Surroundings (accessibility)
- Infrastructure



Discussion


Exchange between countries: How is quality of life for persons with SCI in?

-Which environmental factors play a role?-Which health care factors play a role?-Are there any other factors which play a role?





Life satisfaction and life events



Lucas RE. Long-term disability is associated with lasting changes in subjective well-being: evidence from two nationally representative longitudinal studies. J Pers Soc Psychol 2007;92:717-730.





But what is quality of life?

Quality of life (subjective)

Cognitive evaluation: Life satisfaction Emotional evaluation: Emotional well-being or mental health




Spinal cord injury and Quality of Life

"It is so difficult to accept this terrible life"



"I found out that also with my SCI I can be succesful, independent, handsome, cheerful, and happy"










Measurement occasions

Measurement	N	When?
T1	225	Start of active rehabilitation
Т2	155	3 months after start
Т3	198	Discharge from rehabilitation
Τ4	156	1 year after discharge
Т5	99	2 years after discharge
Т6	146	5 years after discharge



- What is **the course of** life satisfaction of persons with SCI during and after initial rehabilitation up to five years after discharge?
- What is the influence of demographics, SCI characteristics, physical factors, and psychosocial factors on life satisfaction of persons with SCI during and after initial rehabilitation?



Course of life satisfaction after spinal cord injury

1. What is your quality of life at the moment? (1= very unsatisfying, 6= very satisfying)

2. Is your quality of life at the moment worse, equal or better than before the SCI? (1= much worse, 7= much better)

→ Sum score of both questions (range 2-13)



Course of life satisfaction after spinal cord injury

What do you expect?



Methods


Latent Class Growth Mixture Model


Based on regression- and structural equation models.

Muthén LK, Muthén BO. Mplus User's Guide. Sixth Edition. Los Angeles, CA: Muthén & Muthén 1998-2010:197-232.



Five life satisfaction trajectories








Main findings

- Five different life satisfaction trajectories
- Two risk groups: low life satisfaction, deteriorating
- Monitoring during rehabilitation can help to identify persons at risk for poor long-term adjustment



Determinants of quality of life: ICF model





Determinants

- Functional impairment
- Pain and secondary complications
- Social support (emotional support, practical support and esteem support)
- Personality: self-efficacy, appraisals (helplessness, acceptance)



Clinical implications

- **Maximize** functional independence
- Minimize secondary impairments and pain
- **Psychological screening** at the start of active rehabilitation and 3 months after: Identify persons at risk
- Develop psychosocial interventions in persons with SCI

The Dutch SCI rehabilitation: psychosocial



Three phases

- Hospital phase
- Rehabilitation phase
- Chronic phase





NVDG

The <u>Dutch-Flemish Society of Paraplegia</u> exists since 1993 as a society of co-workers from the specialized units for SCI rehabilitation. Goal is to stimulate:

- The collaboration between persons working in the rehabilitation of SCI;
- The exchange of ideas and experience between these persons (national and international);
- Research within the field of SCI rehabilitation.



Patient organisation

• Dwarslaesie Organisatie Nederland

www.dwarslaesie.nl



Psychosocial rehabilitation: Who?

Standard disciplines:

- Intake psychologist
- Intake social worker

Standard or optional:

- Intake peer counselor
- Intake nurse specialised in sexuality
- Intake creative therapist
- Intake sport therapist
- Intake humanistic worker

Consultation:

- Psychiatrist
- Sexuologist



Psychosocial screening

- Adjustment: depression, anxiety, posttraumatic stress
- Personality: strengths and weaknesses
- Coping and appraisals
- Cognition: planning and organisation
- Social network

Outcome:

Inclusion in one of three psychosocial profiles: Resilient, recovery, low life satisfaction



Treatment: Resilient group

- Education about SCI and about adjustment
- Main focus is on physical rehabilitation



Treatment: Recovery group

- Education about SCI and adjustment
- Physical and psychosocial rehabilitation
- Psychosocial treatment involves: EMDR (trauma) Graded activity (depression) Graded exposure (anxiety) Acceptance and commitment therapy Psychomotor therapy (body image) System therapy (partner, children)



Treatment: Low life satisfaction group

- Education about SCI and adjustment
- Main focus is on psychosocial rehabilitation:

Team counseling: how to colloborate together?

Focus on **self-management**: what is possible and what is not possible?

Arrange support in society: family doctor, home care, peer support, keep a link with the rehabilitation center \rightarrow transmural nurse





Power study



Decreased length inpatient care


High level of burden carers

Felt gap after discharge clinic






Aims POWER



PUWER STUDY







Intervention Family Group Conference

	When
Start	End of inpatient rehab.
Future	1,5 months after discharge
End	3 months after discharge
LING	e monthe arter alconarge



- 3 meetings led by a social worker
- Involving social network members (neighbours, family, friends) and professionals
- Together make concrete participation goals and an action plan



Further research in the Netherlands

- Peer support: How is it organised in every SCI rehabilitation center in the Netherlands and what works best?
- Partners for partners organised by the patient organisation



Discussion

- How can patient organisations and rehabilitation centers work together?
- How do patient organisations try to reach persons with SCI?



Case description: resilient



Case description: recovery



Case description: low life satisfaction

De Hoogstraat

ESCIF

- "to live a more independent and full life and participate as any other citizen"
- "to develop your own life project based on your needs and objectives"
- "take charge of your own life instead of having others doing it"



ESCIF: services regarding QOL

- Sports and leisure time activities
- Information
- Legislation and policy lobbying
- Advocacy
- Individual counselling
- Advice and information on health issues
- Peer support
- Accessible accommodations and transportation
- Vocational training
- Skills for independent living
- Courses, Camps, Seminars, support service



ESCIF: self-motivation, selfdetermination, empowerment, coaching and/or peer-support

- Social and vocational rehabilitation
- Personal assistance
- Person centered goal setting
- Self-help groups
- Self-advocacy courses.



What can patient organisations do?

- Make contact during clinical rehabilitation. Lower the threshold by giving education (technical aids, healthy lifestyle), organising activities, facilitating peer support.
- Organise social activities during the weekends in inpatient rehabilitation, such as sport games, going to a theatre/restaurant.
- There is a need for role models. Share your story, also with professionals: videos, presentations, books
- Set up a peer support system for persons in the recovery trajectory and low life satisfaction trajectory and discuss with the person with SCI and the rehabilitation team what the focus should be of everyone who is involved (see Power study).



ESCIF SIP QOL

List of facilitators and barriers

Acceptance from the public	Consumer organisations to enforce the rights and interests of people living with SCI	Peers to exchange experiences
Accessibility	Consumer organisations to help people to get back to life	Professional and leisure time activities
Access to information	Knowledge about SCI to avoid complications	Technical aids – advice and help with the choice, finances, maintenance
Adequate medical care and rehabilitation	Legal support	